

# Using Medications: Why and How?

## Part B: What Should We Know When Taking Medications?



### Prescription (Rx) Label:

**Rx** **Local Pharmacy**  
123 Main St.  
Corner Brook, NL  
709-555-1111


Rx #: 171292                      Date: 25-Jan-2024  
Charlie Bryant

**TAKE 1 TABLET BY MOUTH ONCE DAILY IN AM**  
**CONCERTA 18MG TABLETS**

Keep Out Of Reach Of Children



 **MORNING**

 **SWALLOW WHOLE**  
DO NOT Crush,  
Break, or Chew.

 **MEDICATION SHOULD**  
BE TAKEN WITH PLENTY OF  
**WATER**

# Using Medications: Why and How?

## Part B: What Should We Know When Taking Medications?



### Prescription (Rx) Label:

**Rx** **Local Pharmacy**  
123 Main St.  
Happy Valley-Goose Bay, NL  
709-555-2222

Rx #: 204991                      Date: 4-Feb-2024  
Tamara Bing

**USE ONE SPRAY IN EACH NOSTRIL ONCE DAILY**  
**MOMETASONE NASAL SPRAY (50MCG/SPRAY)**

Keep Out Of Reach Of Children



 **SHAKE WELL**

 **FOR THE *NOSE***

**STORE AT ROOM TEMPERATURE**

**DO NOT SWALLOW**

# Using Medications: Why and How?

## Part B: What Should We Know When Taking Medications?

### Prescription (Rx) Label:


**Rx** **Local Pharmacy**  
123 Main St.  
Lewisporte, NL  
709-555-3333

Rx #: 212332                      Date: 29-Mar-2024  
Lola Martinez

**Use as directed**

**EPIPEN JR    0.15 MG/0.3ML**

Keep Out Of Reach Of Children



**WARNING:** The needle comes out of the orange tip. Do not put your thumb, fingers or hand over the orange tip. DO NOT remove the blue safety release until ready to use.

**STORE AT ROOM  
TEMPERATURE**

# Using Medications: Why and How?



## Part B: What Should We Know When Taking Medications?

### Prescription (Rx) Label:

**Rx** **Local Pharmacy**  
123 Main St.  
Burin, NL  
709-555-4444

Rx #: 109821                      Date: 18-Feb-2024  
Jamal Abdul  
**INHALE 2 PUFFS TWICE DAILY**  
**FLOVENT INHALER 100MCG/PUFF**

Keep Out Of Reach Of Children



# Using Medications: Why and How?

## Part B: What Should We Know When Taking Medications?




### Prescription (Rx) Label:

**Rx** Local Pharmacy  
123 Main St.  
Conception Bay South, NL  
709-555-5555

Rx #: 200128 Date: 12-April-2024  
Jamie Landry

**TAKE 5ML THREE TIMES DAILY FOR 10 DAYS**  
**AMOXICILLIN 250MG/5ML**

Keep Out Of Reach Of Children





**SHAKE WELL**  
STORE IN REFRIGERATOR AND DISCARD  
UNUSED PORTION AFTER 14 DAYS.

# Using Medications: Why and How?

## Part B: What Should We Know When Taking Medications?

### Prescription (Rx) Label:

#### Front of box:



#### Back of box:

<b>Drug Facts</b>
<b>Active ingredients</b> (in each gram) Gramicidin 0.26 mg (Antibiotic) Polymyxin B 0.1 mg (Antibiotic) Lidocaine HCl 45 mg (Analgesic)
<b>Warnings</b> <b>For external use only</b> <b>Do not use</b> : in the eyes, if allergic to any of the ingredients <b>When using this product</b> avoid contact with eyes. If contact occurs, rinse with water <b>Keep out of reach of children</b>
<b>Directions</b> -For adults and children 2 years and older, - for children 2-12 years, application should be supervised by an adult, - clean affected area and apply a small amount of this product on the area 1 to 3 times per day
<b>Other information</b> • store at 15 to 30°C



# Using Medications: Why and How?

## Part B: What Should We Know When Taking Medications?

### Prescription (Rx) Label:

#### Front of box:



#### Back of box:

<b>Drug Facts</b>
<b>Active ingredients</b> (in each tablet) Dimenhydrinate 50 mg (Analgesic)
<b>Warnings</b> Do not give to children under 2 years of age Keep out of reach of children
<b>Directions</b> <ul style="list-style-type: none"><li>• To prevent motion sickness, the first dose should be taken ½ to 1 hour before starting activity (travel in vehicle, boat, airplane)</li><li>• Adults: take 1 to 2 tablets every 4-6 hours, do not take more than 8 tablets in 24 hours, or as directed by a doctor</li><li>• Children (6-12 years old): give 1/2 to 1 tablet every 6-8 hours, do not give more than 3 tablets in 24 hours, or as directed by a doctor</li><li>• Children (2-6 years old): give 1/4 to 1/2 tablet every 6-8 hours, do not give more than 1-1/2 tablets in 24 hours, or as directed by a doctor</li></ul>



# Using Medications: Why and How?

## Part B: What Should We Know When Taking Medications?

### Prescription (Rx) Label:

**Children's Ibuprofen Suspension**  
Ibuprofen 100 mg/5 mL  
Oral Suspension USP

- Relieves pain
- Reduces fever

**For children aged 2-12**

**GRAPE FLAVOUR**

250 ml

**Children's Ibuprofen Suspension**  
Ibuprofen 100 mg/5 mL  
Oral Suspension USP

- Relieves pain
- Reduces fever

**For children aged 2-12** 250 ml

Drug Facts		
<b>Active ingredients</b> Ibuprofen (Pain Reliever, Fever Reducer)		
<b>Directions</b> -Shake well before use, - May take medicine every 8 hours, - Do not use for more than 5 days		
<b>Warnings</b> <b>Do not use</b> : if allergic to any of the ingredients <b>Keep out of reach of children</b>		
Dose		
Age	Weight	Dose
2-3	24-35 lb	6 ml
4-5	36-51 lb	12 ml
6-8	52-71 lb	16 ml
9-10	72- 89 lb	20 ml
11-12	90-100 lb	22 ml
<b>Other information</b> • store at 15 to 30°C		

**TAKE WITH FOOD**

**SHAKE WELL BEFORE USING**